

5 Davis Road East Old Lyme, CT 06371 Tel 860.390.6000 Fax 860-215-8150

info@northeastpsc.com www.northeastpsc.com

Nilay Shah, MD Megan Phillips, PA-C Vinod V. Pathy, MD, FACS

Jillian Fortier, MD Morgan Massey, PA-C

## Nipple Areola Tattooing

Pre-operative Instructions

ightharpoons	Your scheduled	procedure	date is	

- If you smoke or use any other nicotine products (including the patch or gum), please stop at least 4 weeks prior to surgery. You should refrain from nicotine products for at least 4 weeks following your surgery as well. Smoking reduces circulation and can therefore result in wound healing impairment. It may also increase your risk of capsular contracture formation.
- Avoid the following anti-platelet agents for 7-10 days prior to your surgery, including, but not limited, to: ibuprofen products (Advil, Motrin), Aleve, Naprosyn, Celebrex, Toradol, fish oil, herbals, and supplements including those containing Vitamin E, and generally for 1 week following your surgery, as they result in a higher risk of bleeding.
- If you are taking the following blood thinners or anti-platelet agents, please notify your surgeon AND contact the provider who is managing such medications as soon as possible to obtain instructions if and when you should stop the medication prior to surgery, and which alternative medications may be used in the interim: aspirin, Coumadin (warfarin), Lovenox, Plavix, Pletal, Effient, Aggrenox, Pradaxa, Savaysa, Xerelto, Eliquis, or Brilinta.
- Avoid alcohol for at least 48 hours prior to your procedure.
- Please purchase Bacitracin or Neosporin ointment, nonadherent gauze sponges, and 1" surgical paper tape from your pharmacy prior to your surgery, as you will need them for post-procedure dressing care.
- ➤ You may shower the morning of your surgery. Do not apply any moisturizers, makeup, deodorant, or perfume.
- Wear a large, front closure shirt to the office on the day of your procedure.