



NORTHEAST PLASTIC SURGERY  
CENTER

5 Davis Road East  
Old Lyme, CT 06371

info@northeastpsc.com  
www.northeastpsc.com

Tel 860.390.6000  
Fax 860-215-8150

**Tomer Avraham, MD, FACS**

**Vinod V. Pathy, MD, FACS**

**Charles Karcutskie, MD**

**Megan Phillips, PA-C**

**Hannah Chalmers, PA-C**

## **Gynecomastia Reduction**

### Post-operative Instructions

- ▶ Your post-operative appointment is scheduled for \_\_\_\_\_.
- ▶ Do not remove your dressings. Your dressings will be removed during your first office visit. Your dressings will be removed at your first post-operative visit. Keep your chest garment in place. The garment will be worn for 4-6 weeks. You may place gauze sponges inside your garment for added comfort.
- ▶ Keep your dressings dry. Sponge-bathe only until further instructed. Usually, you may shower 48 hours after final drain removal.
- ▶ Empty your drains every 12 hours and record the measurements for each drain, separately and labeled by side, on your chart. Please bring the chart with you to your office visits. The amounts that are draining from each drain will determine when each drain may be removed. The drains are usually removed within 1 week to 10 days, but a number of factors can influence the duration. Once each drain is removed, you should keep a dry gauze dressing for 48 hours over the former sites on the skin to prevent drainage onto your clothing but then may discontinue the drain dressings altogether after those 48 hours.
- ▶ If a liquid skin adhesive (glue) was used along your incisions, it may take several weeks to dissolve. Do not apply ointments such as Bacitracin directly over the glued area unless instructed, as this may prematurely dissolve this dressing.
- ▶ You have received prescriptions for \_\_\_\_\_ as an antibiotic and \_\_\_\_\_ for pain relief. Postoperative pain management includes other modalities that are non-narcotic based to minimize reliance on such medications. You may stop taking your antibiotic after the final drain has been removed. Please take Colace as a stool softener to limit constipation.
- ▶ Avoid the following anti-platelet agents until at least your first post-op appointment, including, but not limited, to: ibuprofen products (Advil, Motrin), Aleve, Naprosyn, Celebrex, Toradol, fish oil, herbals, and supplements including those containing Vitamin E, as they result in a higher risk of bleeding.
- ▶ If you are taking the following blood thinners or anti-platelet agents, you should have already obtained instructions pre-operatively when they may resume, but please discuss with your surgical team: aspirin, Coumadin (warfarin), Lovenox, Plavix, Pletal, Effient, Aggrenox, Pradaxa, Savaysa, Xarelto, Eliquis, or Brilinta.

- ▶ Do not consume alcohol while you are taking any medications following surgery, especially pain medications, muscle relaxants, anti-anxiety medications, sleeping pills, or other sedatives.
- ▶ You may resume your pre-operative diet as tolerated.
- ▶ Restrictions are as follows:
  - ▶ Do not lay flat in bed. Sleep with your head elevated on at least 2 pillows for 2 weeks. Do not lay on your sides for 6 weeks.
  - ▶ You may walk as tolerated beginning the day of your surgery.
  - ▶ No driving for 2 weeks. You will be restricted from driving until all narcotic and muscle relaxant medications have been stopped.
  - ▶ No heavy lifting (> 5 lbs) for 4 weeks.
  - ▶ No strenuous pulling or pushing for 4 weeks.
  - ▶ No vigorous upper body workout for 6 weeks.
  - ▶ Avoid hot or cold packs against your skin, as skin changes may occur because of sensory changes.
- ▶ Please refrain from nicotine-containing products for at least 4 weeks following your surgery. Nicotine reduces circulation and can therefore result in wound healing impairment.
- ▶ Based upon the mechanical nature of the procedure, fat grafting can often result in bruising at the donor fat harvest site. In the majority of patients, this bruising resolves in 1-3 weeks.
- ▶ Every patient and procedure is unique. However, please alert us if you observe any of the following and are concerned:
  - ▷ temperature  $\geq 101.4^{\circ}\text{F}$
  - ▷ increased redness at the surgical site or adjacent site(s)
  - ▷ worsening bruising at the surgical or adjacent site(s)
  - ▷ blisters at the surgical site(s)
  - ▷ increased swelling or size leading to significant and sudden asymmetry
  - ▷ unremitting or increasing pain at the surgical site(s)
  - ▷ significant drainage at the surgical site
- ▶ There are a number of different modalities that can limit scar formation post-operatively. Starting 2 weeks post-operatively, we may recommend that you utilize silicone sheeting such as Epi-Derm™ Silicone Gel Strips from Biodermis™. This therapy may be purchased directly through [www.biodermis.com](http://www.biodermis.com). Silicone sheets should be applied overlying the healing scar for at least 12 hours daily, and ideally closer to 24 hours daily, with gentle cleansing of the scar performed once daily. An individual sheet may be re-used multiple times, with the average lifespan being 7-10 days. Alternatively, you may choose to utilize any number of silicone scar reduction gels, such as Mederma® or Biocorneum®, and massage in a thin coat twice daily to the scar for 2-3 months. We recommend silicone gel with the added embedded SPF component.