

5 Davis Road East Old Lyme, CT 06371

info@northeastpsc.com www.northeastpsc.com Tel 860.390.6000 Fax 860-215-8150

Vinod V. Pathy, MD, FACS

Charles Karcutskie, MD

Megan Phillips, PA-C

Hannah Chalmers, PA-C

## **Brazilian Butt Lift**

Pre-operative Instructions

	Your scheduled surgery date is		
<b>&gt;</b>	Your procedure will be performed at	·	
<b>&gt;</b>	You should call Surgical Booking at day prior to your scheduled procedure to determine as to when to facility.		
<b>&gt;</b>	Labwork has been ordered for you in preparation for your surgery, which may include an EKG and/or a chest X-ray as well. Please complete all tests at least 2 weeks prior to your procedure, but no more than 4 weeks before your scheduled surgical date.		
<b>&gt;</b>	f you smoke or use any other nicotine products (including the patch or gum), please stop at least 6 weeks prior to surgery. You should refrain from nicotine products for at east 6 weeks following your surgery as well. Smoking reduces circulation and can therefore result in wound healing impairment and decreased fat grafting survival.		
<b>&gt;</b>	Avoid alcohol for at least 48 hours prior to your surgery.		
<b>&gt;</b>	Avoid the following anti-platelet agents for 7-10 days prior to your subut not limited, to: ibuprofen products (Advil, Motrin), Aleve, Nap Toradol, fish oil, herbals, and supplements including those containing Name to the following your surgery, as they result in a higher risk of bleeding.	orosyn, Celebrex, itamin E, and for	
<b>&gt;</b>	If taking the following blood thinners or anti-platelet agents, please notify your surgeon AND contact the provider managing such medications as soon as possible to obtain instructions if and when you should stop the medication prior to surgery, and which alternative medications may be used in the interim: aspirin, Coumadin (warfarin), Lovenox, Plavix, Pletal, Effient, Aggrenox, Pradaxa, Savaysa, Xerelto, Eliquis, or Brilinta.		
<b>&gt;</b>	You have received a prescription for as antibiotic and for post-operative pain relie take <u>Colace</u> post-operatively to limit constipation. Please fill these pre 1 week prior to your surgical date.	a post-operative f. Please plan to scriptions at least	

- ▶ Bring all of your prescribed and regularly used over-the-counter medications with their original bottles to the facility on the day of surgery.
- Please purchase 4"x4" gauze and 1" surgical paper tape from your pharmacy prior to your surgery, as you may need them for post-operative dressing care.
- ▶ We will assist in obtaining a suitable compression garment for usage post-operatively which you will bring to the operating room on the day of your procedure. Often, patients choose to purchase a second garment for ease of cleaning and wear.
- Do not eat or drink anything starting at midnight on the night before your surgery. If discussed at your pre-op visit, you may be able to take selected medications with a sip of water on the morning of your surgery. Please refer to the additional instruction sheet if you are on medical weight loss treatment.
- You may shower the morning of your surgery. Do not apply any moisturizers, makeup, deodorant, or perfume.
- Remove all jewelry, rings, necklaces, and piercings, prior to your operating room visit. Leave all non-essential valuables at home.
- ▶ Avoid wearing contact lenses to your operating room visit.
- ▶ Wear a large, front closure shirt and loose elastic waisted pants to the hospital on the day of your surgery.
- Arrange for appropriate transportation in advance of your surgery. You are required to have a family member or friend bring you to and from the facility, before and after your surgery, as you will be unable to safely transport yourself home post-operatively.