



NORTHEAST PLASTIC SURGERY
CENTER

5 Davis Road East
Old Lyme, CT 06371

info@northeastpsc.com
www.northeastpsc.com

Tel 860.390.6000
Fax 860-215-8150

Vinod V. Pathy, MD, FACS

Megan Phillips, PA-C

Charles Karcutskie, MD

Hannah Urgo, PA-C

Blepharoplasty

Post-operative Instructions

- ▶ Your post-operative appointment is scheduled for _____.
- ▶ Keep your head elevated on at least 2 pillows for at least the next 72 hours. This will help limit the swelling that you will incur at your eyelids. Sleep on your back and not on either side.
- ▶ Apply cold, moist gauze to your eyelids to act as cool compresses for at least the next 72 hours. These may be fashioned by dipping 4"x4" gauze sponges in a slush of crushed ice before applying to the eyelids. Do not apply ice packs to your eyelids, as the excess weight may be uncomfortable and may impair recovery.
- ▶ Apply artificial tear drops to your eyes as needed to avoid dry eyes.
- ▶ You may shower 24 hours after the surgery. Do not rub or scrub the eyelids. Simply allow soap and water to run over the sites and pat dry afterwards. Do not attempt to get rid of all the crusting or dried blood, as this can disrupt the closure.
- ▶ You have received prescriptions for _____ for pain relief, however, this can typically be avoided. Postoperative pain management includes other modalities that are non-narcotic based (including acetaminophen or Tylenol) to minimize reliance on such medications. Please take Colace as a stool softener to limit constipation.
- ▶ Avoid the following anti-platelet agents until 24 hours after your surgery, including, but not limited, to: ibuprofen products (Advil, Motrin), Aleve, Naprosyn, Celebrex, Toradol. You may also resume any of the following supplements after 24 hours: fish oil, herbals, and supplements including those containing Vitamin E.
- ▶ If you are taking the following blood thinners or anti-platelet agents, you should have already obtained instructions pre-operatively when they may resume, but please discuss with your surgical team: aspirin, Coumadin (warfarin), Lovenox, Plavix, Pletal, Effient, Aggrenox, Pradaxa, Savaysa, Xerelto, Eliquis, or Brilinta.
- ▶ Do not consume alcohol while you are taking any medications following surgery, especially pain medications, muscle relaxants, anti-anxiety medications, sleeping pills, or other sedatives.

- ▶ You may resume your pre-operative diet as tolerated.
- ▶ Restrictions are as follows:
 - ▶ Reading may resume after 2-3 days.
 - ▶ Avoid contact lenses for 2 weeks.
 - ▶ Avoid strenuous activity for 3 weeks.
 - ▶ Avoid leaning over with your head down for 3 weeks.
 - ▶ Avoid alcohol for 3 weeks.
- ▶ Please refrain from nicotine-containing products for at least 6 weeks following your surgery. Nicotine reduces circulation and can therefore result in wound healing impairment.
- ▶ Every patient and procedure is unique. However, please alert us if you observe any of the following and are concerned:
 - ▷ temperature $\geq 101.4^{\circ}\text{F}$
 - ▷ increased redness at the surgical site or adjacent site(s)
 - ▷ worsening bruising at the surgical or adjacent site(s)
 - ▷ blisters at the surgical site(s)
 - ▷ increased swelling or size leading to significant and sudden asymmetry
 - ▷ unremitting or increasing pain at the surgical site(s)
 - ▷ significant drainage at the surgical site